

Piecewise Monotone Approximation of Unbounded Functions In Weighted Space $L_{p,w}([-\pi,\pi])$



Alaa Adnan Auad^{1*} and Sultan Hameed Mehiady²

¹Department of Mathematics, College of Education for pure science, University of Anbar, Ramadi, Iraq

²Department of Mathematics, College of Education for pure science, University of Anbar, Ramadi, Iraq

ARTICLE INFO

Received: 4 / 4 / 2022
 Accepted: 16 / 5 / 2022
 Available online: 20/7/2022

DOI: 10.37652/juaps.2022.174841

Keywords:

piecewise monotone approximation.
 unbounded functions.
 weight space.

ABSTRACT

In this paper, investigate the approximation of unbounded functions in weighted space, by using trigonometric polynomials considered. We introduced type of polynomials piecewise monotone (g_{κ}) having same local monotonicity as unbounded functions without affecting the order of huge error have a finite number of max. and min. unbounded functions that amount. In addition, we established not included any of extreme points of this functions, χ of and closed subset γ on closed intervals χ then there exist class of polynomials such that the best of approximation has high or order of β and such that for κ sufficiently great of the polynomials and functions have the same monotonicity at each of γ .

1. INTRODUCTION

Let $\chi = [-\pi, \pi]$ be closed intervals, $L_p([-\pi, \pi])$ we can be define as the space of all bounded function on χ , with the norms.

$$\|f\|_p = \left(\int_{\chi} |f(t)|^p dt \right)^{\frac{1}{p}} < \infty$$

Let $\mathcal{C}(w)$ be the set of all weighted function on χ , a weighted function can be written. The follows $w : \chi \rightarrow \mathcal{R}^+$ such that $|f(t)| \leq M/w(t)$, $w > 0$, $M \in \mathcal{R}^+$, $w(t) \in \mathcal{C}(w)$ and $L_{p,w}([-\pi, \pi])$, is the space of all unbounded functions on χ , which are equipped the norms.

$$\|f\|_{p,w} = \left(\int_{\chi} |f(t).w(t)|^p dt \right)^{\frac{1}{p}} < \infty$$

Let \mathbb{T}_{κ} be the set Which contains any polynomials of degree less than or equal to κ .

Let $[a, b]$ be a compact interval and $f : [a, b] \rightarrow \mathcal{R}^+$ a function, f is piecewise monotone if and only if there exists a partition $a = t_1 < t_2 < t_3 < \dots < t_{\kappa} = b$, on $[a, b]$ and a, b will be called the topes of f , such that $f_i = f|_{(t_{i-1}, t_i)}$ is monotone for all $i = 1, 2, \dots, \kappa - 1$.

The aim of this paper is discuss piecewise monotone approximation of unbounded functions in weighted space

*Corresponding author at Department of Mathematics/College of Education for pure science/University of Anbar, E-mail address: alaa.adnan.auad@uoanbar.edu.iq

by some types of efficient polynomials. An example [1], [2] and [3]. The strongest results are by [4], [5], [6] and [7]. In 2011, kopotun [8], introduce the point wise shape which preserving approximation of function by algebraic polynomials. In fact Dzyubenko [9] and Leviatan [10] have obtain interpolatory estimates in monotone piecewise polynomial approximation and both of them shown good result in monotone approximation.

So, we can define the operator as:

$$G_{\kappa}(t) = \frac{1}{g_{\kappa}} \left(\frac{\sin \kappa t / 2}{\sin t / 2} \right)^4; \text{ Where } g_{\kappa} = \int_{-\pi}^{\pi} \left(\frac{\sin \kappa t / 2}{\sin t / 2} \right)^4 dt.$$

$$\text{And } \kappa^3 \leq g_{\kappa} \leq 2\pi \kappa^3.$$

2. Auxiliary Lemmas

Lemma 2.1. The operator $G_{\kappa}(t)$ has the following properties:

(i) If κ is even, $0 < \beta \leq \pi$, then

$$\int_{-\pi}^{\pi} G_{\kappa}(t) dt \leq \int_0^{\beta} G_{\kappa}(t) dt.$$

(ii) if, $0 < \beta \leq \pi/2$, then

$$\int_{\beta}^{\pi} G_{\kappa}(t) dt \leq \frac{C}{\kappa^3 \beta^3}.$$

Where C is positive constant of real numbers.

Proof: (i) We take $0 < \beta \leq \pi/2$;

$G_\kappa(t)$ is an even periodic and we have

$$\int_{-\pi}^{\pi+\beta} G_\kappa(t) dt = \int_{\pi-\beta}^{\pi} G_\kappa(t) dt .$$

$$\text{So , } \int_0^\beta [G_\kappa(t) - G_\kappa(t + \pi)] dt = \int_0^\beta G_\kappa(t) dt -$$

$$\int_{\pi}^{\pi+\beta} G_\kappa(t) dt \quad \dots (2.1)$$

Also, if $\pi/2 < \beta \leq \pi$, Then

$$\int_0^\beta G_\kappa(t) dt - \int_{\pi-\beta}^{\pi} G_\kappa(t) dt = \int_0^{\pi-\beta} G_\kappa(t) dt -$$

$$\int_{\beta}^{\pi} G_\kappa(t) dt \quad \dots (2.2)$$

From (2.1) and (2.2) we obtain

$$\int_{-\pi}^{\pi} G_\kappa(t) dt \leq \int_0^\beta G_\kappa(t) dt .$$

(ii) Let $0 < \beta \leq \pi/2$. Then, since $\kappa^3 \leq g_\kappa \leq 2\pi\kappa^3$.

$$\text{So, } \int_{\beta}^{\pi} G_\kappa(t) dt \leq \frac{1}{\kappa^3} \int_{\beta}^{\pi} \left(\frac{\sin \kappa t/2}{\sin t/2}\right)^4 dt \leq \frac{1}{\kappa^3} \int_{\beta}^{\pi} \frac{1}{(\sin t/2)^4} dt .$$

Now, for $0 < t \leq \pi$. $\sin t/2 \geq t/2$, hence

$$\int_{\beta}^{\pi} G_\kappa(t) dt \leq \frac{\pi^4}{\kappa^3} \int_{\beta}^{\pi} \frac{1}{t^4} dt = \frac{\pi^4}{3\kappa^3} \left[\frac{-1}{\pi^4} + \frac{1}{\beta^4}\right] \leq \frac{C}{\kappa^3 \beta^3}$$

And this lemma is proving.

3. Main results

Theorem 3.1 :

Let $t \in [-\pi, \pi]$, $f \in L_{p,w}([-\pi, \pi])$, $1 \leq P < \infty$ Periodic function. Then there exist trigonometric polynomial q_κ of order less than or equal to κ , such that

$$\|f - q_\kappa\|_{p,w} \leq \frac{C}{\kappa} , \text{ where } C \text{ is positive constant.}$$

Proof. we can choose $q_\kappa(t) = q_\kappa(t + 2\kappa\pi)$, $\kappa = 1, 2, \dots$ and

$$q_\kappa(t) = \int_{-\pi}^{\pi} G_\kappa(\theta) f(t - \theta) d\theta$$

Since f Periodic function, we have

$$q_\kappa(t) = \int_{-\pi}^{\pi} G_\kappa(\theta) f(t - \theta) d\theta = \int_{t-\pi}^t G_\kappa(\theta) f(t - \theta) d\theta -$$

$$\int_t^{t+\pi} G_\kappa(\theta) f(t - \theta) d\theta$$

So,

$$q'_\kappa(t) = \int_{t-\pi}^t G_\kappa(\theta) d\theta - \pi f(t - \pi) - \int_t^{t+\pi} G_\kappa(\theta) d\theta -$$

$$\pi f(t + \pi)$$

$$= \int_{t-\pi}^t G_\kappa(\theta) d\theta - \int_t^{t+\pi} G_\kappa(\theta) d\theta$$

We can take $0 < t \leq \pi$, and $G_\kappa(\theta)$ is an periodicity. From

Lemma 2.1, we obtain

$$q'_\kappa(t) = 2 \int_0^t G_\kappa(\theta) d\theta - 2 \int_{\pi-t}^{\pi} G_\kappa(\theta) d\theta$$

$$\text{Since, } 2 \int_0^t G_\kappa(\theta) d\theta \geq 2 \int_{\pi-t}^{\pi} G_\kappa(\theta) d\theta$$

$$\text{Implies } q'_\kappa(t) \geq 0$$

Also, for $-\pi < t \leq 0$, we obtained $q'_\kappa(t) \leq 0$, $q_\kappa(t)$ non-negative and,

$$|q_\kappa - f| \geq 0, \quad \text{implies } \|q_\kappa - f\|_{p,w} \leq \frac{C}{\kappa} .$$

Theorem 3.2:

Let $q_\kappa(t)$ be a monotone function in $L_{p,w} \left(\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \right)$. Then there exist a piecewise linear operator F , such that

$$\|q_\kappa - F\|_{p,w} \leq \frac{C}{\kappa} \text{Max}(F), \text{ where } C \text{ is positive}$$

constant.

Proof. Let $-\frac{\pi}{2} = t_1 < t_2 < t_3 < \dots < t_k = \frac{\pi}{2}$

We need to make the $q_\kappa \in \mathbb{T}_k$, such that satisfying

$$\|q_\kappa - F\|_{p,w} \leq \frac{C}{\kappa} \text{Max}(F)$$

$$\text{Let } \theta_0 = \frac{p_0 + p_{\kappa-1}}{2}, \theta_i = \frac{p_i - p_{i-1}}{2}; \text{ For } i = 1, 2, \dots, \kappa - 1.$$

Thus,

$$F(t) = K +$$

$$\sum_{i=0}^{\kappa-1} \theta_i |t - s_i|$$

$$\dots (3.1)$$

Where $-\frac{\pi}{2} = s_1 < s_2 < s_3 < \dots < s_k = \frac{\pi}{2}$ and K is

constant.

Let q be defined by theorem (1) and, let

$$q_\kappa(t) = K +$$

$$\sum_{i=0}^{\kappa-1} \theta_i q(t - s_i)$$

$$\dots (3.2)$$

Where $q_\kappa \in \mathbb{T}_k$. So, from (3.1), and (3.2), we get

$$\|q_\kappa - F\|_{p,w} \leq \frac{C}{\kappa} \text{Max} \left| \sum_{i=0}^{\kappa-1} \theta_i \right| \leq \frac{C}{\kappa} \text{Max}(p_i) =$$

$$\frac{C}{\kappa} \text{Max}(F) .$$

Theorem 3.3:

Let $f \in L_{p,w} \left(\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \right)$, $1 \leq P < \infty$ and C is positive constant independent of k . Then there exist a monotone function $\langle q_\kappa \rangle$, $q_\kappa \in \mathbb{T}_k$ such that

$$\|f - q_\kappa\|_{p,w} \leq \frac{\text{Max}(C)}{\kappa} .$$

Where f is the proper piecewise monotone functions.

Proof. Let F_k be the proper piecewise linear operator define

on $\left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$, such that its has nodes at the topes of the function f

at the points $-\frac{\pi}{2} + \frac{i\pi}{\kappa}$, for $i = 0, 1, 2, \dots, \kappa$, and $F_{\kappa}(t) = f(t)$, since $\|F_{\kappa} - f\|_{p,w} \leq \frac{C_1}{\kappa}$,

where C is positive constant and $Max(F) \leq \frac{C_1}{\kappa}$. By using theorem (2), there is a Polynomial $q_{\kappa} \in \mathbb{T}_{\kappa}$ such that

$$\|F_{\kappa} - q_{\kappa}\|_{p,w} \leq \frac{C_2}{\kappa}.$$

We need to prove that

$$\|f - q_{\kappa}\|_{p,w} \leq \frac{Max(C)}{\kappa}.$$

So,

$$\begin{aligned} \|f - q_{\kappa}\|_{p,w} &= \left(\int_X |f(t) - q_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} \\ &= \left(\int_X |f(t) - F_{\kappa}(t) + F_{\kappa}(t) - q_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} \\ &\leq \left(\int_X |f(t) - F_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} + \left(\int_X |F_{\kappa}(t) - q_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} \\ &\leq Max \left[\left(\int_X |f(t) - F_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} + \left(\int_X |F_{\kappa}(t) - q_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} \right] \\ &\leq Max \left(\int_X |f(t) - F_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} + Max \left(\int_X |F_{\kappa}(t) - q_{\kappa}(t)|^p w(t) dt \right)^{\frac{1}{p}} \\ &\leq Max(C) \|F_{\kappa} - f\|_{p,w} + Max(C) \|F_{\kappa} - q_{\kappa}\|_{p,w} \\ &\leq \frac{Max(C_1)}{\kappa} + \frac{Max(C_2)}{\kappa} \\ &\leq \frac{Max(C)}{\kappa}. \end{aligned}$$

Theorem 3.4 :

Let $f \in L_{p,w}([a, b])$, $1 \leq p < \infty$. Then there exist a nearly monotone operator

$\langle \mathcal{P}_{\kappa} \rangle$, $\mathcal{P}_{\kappa} \in \mathbb{P}_{\kappa}$ such that

$$\|\mathcal{P}_{\kappa} - f\|_{p,w} \leq \frac{c}{\kappa} Max(F).$$

Where f is the proper piecewise monotone on $[a, b]$.

Proof: The proof of this theorem by using the operator $q_{\kappa}(t) = \cos(t)$ with some modification. Of theorem 3.3

References

[1] Darst R. B. and sahib s., (1983), Approximation of continuous and quasi - continuous functions by monotone Functions, j. Approx. Theory 38 ,p.p. 9 - 27.

[2] Hectare R. , (1985), Best L1 -approximation of quasi - continuous functions on [0,1] by non-decreasing functions, Functions, j. Approx. Theory 44, p.p. 221 - 229. J. New York. Devore A. and Lorentz G., (1993), Constructive Approximation, Springer-verlag [3]

[4] Leviathan D. , (1998) ,Monotone approximation estimates involving the third modulus of smoothness, Approximation I.(Nashville, TN, 1998), Innov - App..Math. Vanderbilt univ. press, Nashville, TN, ,p Theory IX, NoI..p.233 - 230.

[5] Koptune K.A., (1994) ,Point wise and uniform estimates for convex approximation of functions by Algebraic polynomial , Constar. Approx. to ,p.p.153-178.

[6] Wilbert D., (2004) , Best approximation by the inverse of a monotone polynomial and location problem, J. Approx. Theory (114), 98 - 114.

[7] Alaa A.A. and Alaa M., (2017), co positive approximation of unbounded functions in weighted space,international Of Applied Engineering Research, vol.12 No. 21 , p.p. 11392 - 11396. Journal

[8] Koptune K.A., Leviathan D. and shevechuk, I. A., (2011), uniform and point wise shape preserving approximation by Algebraic polynomials, surv. Approx. Theory 6 24 - 74.

[9] Dzyubenko, G.A., Leviathan D., (2014), point wise estimates of convex approximation, Jaen J. Approx 6 ,P.P. 261 - 295.

[10] Leviathan D. and petrova I. L. ,(2017), interpolatory estimates in monotone piecewise polynomial approximation J. Approx. Theory 233 ,p.p. 1 - 8.

أفضل التقريب الرتيب المتقطع للدوال الغير مقيدة في الفضاء الموزون

علاء عدنان عواد¹ سلطان حميد مهدي²

¹ قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة الأنبار، رمادي، العراق

² قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة الأنبار، رمادي، العراق

الخلاصة:

في هذا البحث سنعرض تقريب الدوال الفضاء الوزون بواسطة متعددات مثلثية ونقدم نوع من المتعددات الرتيبة المتقطعة والتي لها نفس الرتبة الموقعية للدوال الغير مقيدة دون تأثير الخطا الكبير والتي يكون لها العدد المنتهي من الحدود العليا والدنيا اضافة الى ذلك عدم تضمين النقاط المتطرفة لهذه الدوال في الفترات الجزئية المغلقة للفترة الاساسية المغلقة لذا سوف سيتولد لنا فئة من متعددات الحدود من خلالها نحصل على أفضل تقريب من رتبة بيتا يبحث يكون لاي (ك) يكون لهذه الدوال والمتعددات نفس الرتبة.