

Optimized Trigonometric Spline Function with Conjugate Gradient Method for Solving FDEs

Faraidun K.Hamasalh¹, Gulnar W.Sadiq², Emad S.Salam³



¹ Department of Mathematics, College of Education, University of Sulaimani, Sulaimani, Iraq

² Department of Mathematical Sciences, College of Basic Education, University of Sulaimani, Sulaimani, Iraq

³ MSc Student at University of Sulaimani, Sulaimani, Iraq

ARTICLE INFO

Received: 24 /04 / 2022

Accepted: 09 / 08 / 2022

Available online: 22/12/ 2022

DOI: [10.37652/juaps.2022.176477](https://doi.org/10.37652/juaps.2022.176477)

Keywords:

Conjugate gradient,
Non-polynomial spline,
Fractional derivative,
Gauss-Seidel method.

ABSTRACT

We investigate the non-polynomial spline function to solve the fractional differential equations with the conformable conjugate gradient method. The fractional derivative was described using the Caputo fractional derivative to construct the spline scheme with polynomial fractional order. Therefore, transform the problem to an equivalent iterative linear system that can be solved by Gauss-Seidel and conjugate gradient methods. For the given spline function, error bounds were studied and a stability analysis was completed, the error estimation is also calculated as different values of (n) depend on the step size oh (h). Numerical examples with known analytical solutions are shown to verify the method's accuracy. The outcomes are in satisfactory correlation with the exact answers according to the numerical experiments. Moreover, the convergence analysis was investigated with the drive some theorems. Also, the procedure is explained in depth and supported by computational examples and the results show that the fractional spline function which interpolates data is productive and profitable in solving unique problems and compare with the exact solutions.

Introduction:

Due to its numerous applications in science and engineering, fractional calculus plays a vital role in a multitude of fields including. materials modelling[1], electromagnetism[2], processing of signals[3], diffusion processes[4], fluid mechanics[5], electrical engineering[6], mathematical economics[7]. A variety of techniques have been invented to solve fractional differential equations, such as:, fractional finite difference method[8], Adomain decomposition method[9], Adam-Bashforth-Multon method[10],

Homotopy analysis[11], matrix approach method for solving FDE discussed in[12], fractional explicit Adams method used in[13] Muhammad I. Bhatti and Md.Habibur Rahman used B-polynomial bases to solve FDE[14], discrete Prabhakar fractional operator studied in[15], a spectral Tau method investigated by Hari Mohan Srivastava and et al[16].

One of the most effective procedures for solving large linear systems of equations is the conjugate gradient method, which can also be applied to nonlinear optimization. The linear conjugate gradient method was proposed in the 1950s by Hestenes and Stiefel to solve linear systems of equations with positive definite matrices as an alternative to Gauss elimination. [17]. Fletcher and Reeves discussed the non-linear conjugate gradient method in 1964.[18], Several modifications have been introduced by researchers to the CG method, such as (,[19])

Splines have numerous applications in physics and engineering and many researchers have worked on them. For instance, H. Justine and J. Sulaiman constructed a cubic non-polynomial spline for solving BVPs [20], The application of non-polynomial spline to the numerical solution for the fractional differential equation discussed in [21], a six degree spline was found to solve second order initial value problem in [22] .

*Corresponding author at: Department of Mathematics, College of Education, University of Sulaimanli ;Email: faraidun.hamasalh@univsul.edu.iq

In this study we consider the fractional differential equation of the form

$$D^\lambda y + p(x)y' + q(x)y = r(x), \quad x \in [a, b] \quad (1)$$

With boundary conditions

$$y(a) = B_1, \quad y(b) = B_2 \quad (2)$$

Where $p(x), q(x)$ and $r(x)$ are functions of x , B_1 and B_2 are constants. Then the interval $[a, b]$ is uniformly divided into j subintervals and the length of intervals defined as $\Delta x = h = \frac{b-a}{j}, n = j - 1$.

The structure of this paper is organized as follows: we set some basic definitions and outcomes of fractional calculus in section 2, construction of non-polynomial spline function is presented in section 3, error estimations and convergence analysis is in section 4, finally some numerical examples are illustrated in section 5.

2. Mathematical Preliminaries

There are various definitions of fractional derivative and Taylor's theorem, which used in this work, will be presented in this section. The most common definitions of fractional derivatives are Caputo and Riemann-Liouville definitions.

Definition 2.1. [23] The Caputo fractional derivative of order $\lambda > 0$ is defined by

$$D^\lambda f(t) = \frac{1}{\Gamma(m-\lambda)} \int_a^t (t-\tau)^{m-\lambda-1} \frac{d^m}{d\tau^m} f(\tau) d\tau, \\ m-1 < \lambda < m \in N.$$

Definition 2.2. [12] The Riemann-Liouville fractional derivative of order $\lambda > 0$ is defined by

$$D^\lambda f(t) = \frac{1}{\Gamma(m-\lambda)} \frac{d^m}{d\tau^m} \int_a^t (t-\tau)^{m-\lambda-1} f(\tau) d\tau, \\ m-1 < \lambda < m \in N.$$

Definition 2.3.[12] The Riemann-Liouville fractional integral of order $\lambda > 0$ is defined by

$$I^\lambda f(t) = \frac{1}{\Gamma\lambda} \int_a^t (t-\tau)^{\lambda-1} f(\tau) d\tau, \\ m-1 < \lambda < m \in N.$$

Definition 2.4.[14] The Caputo derivative of order λ of a polynomial function x^d is defined by $D^\lambda x^d = \frac{\Gamma(d+1)}{\Gamma(d-\lambda+1)} x^{d-\lambda}$.

Definition 2.5.[24] The Spectral radius $\psi(A)$ where A is an $n \times n$ matrix is given by $\psi(A) = \max(|\lambda|)$ where λ is an eigenvalue of A .

Definition 2.6.[24] An $n \times n$ matrix A is convergence if $\psi(A) < 1$.

Definition 2.7.[25] A square matrix A is called diagonally dominate if

$$|a_{ii}| > \sum_{i \neq j} |a_{ij}|.$$

Theorem 2.1. For any $x_0 \in R^n$ the sequence x_k generated by conjugate gradient method converges to the solution x^* in at most n steps.

Proof. See [17].

3. Analysis of Non-Polynomial Spline Function with Fractional Order

The new model of non-polynomial spline method is to create a grid with step size x_i and fractional order, and also new conditions as the follows:

$$S(x) = S_i(x), x \in [x_i, x_{i+1}], \quad i = 0, 1, 2, \dots, n \quad (3)$$

Here the trigonometric spline function with fractional order by assuming

$$S_i(x) = a_i + b_i(x-x_i)^{\frac{1}{2}} + c_i(x-x_i) + d_i \cos k(x-x_i) + e_i \sin k(x-x_i) \quad (4)$$

Where a_i, b_i, c_i, d_i and e_i are constants for, $i = 0, 1, \dots, n$, and k is a free parameter.

The spline function $S(x)$ interpolates $y(x)$ depending on k . To find the value of constants in (4) we supposed the following conditions

$$S_i(x_i) = y_i, \quad S_i(x_{i+1}) = y_{i+1}, \quad S_i^{(\frac{1}{2})}(x_i) = y_i^{(\frac{1}{2})}, \\ S_i'(x_i) = D_i, \text{ and } S_i'(x_{i+1}) = D_{i+1}. \quad (5)$$

Applying the conditions in equation (5) the value of all constants in (4) obtained as follows

$$a_i = \frac{-\sqrt{\pi} M_1 M_3 y_{i+1} + \sqrt{\pi} M_1 M_3 y_i + 2\sqrt{h} M_1 M_3 y_i^{(\frac{1}{2})} - (\beta M_3 + \alpha) D_i + (\delta M_3 + \alpha) D_{i+1}}{\alpha M_1}, \\ b_i = \frac{y_i^{(\frac{1}{2})} - \sqrt{2k}(\beta M_2 + \alpha) D_i + \sqrt{2k}(\delta M_2 + \alpha) D_{i+1}}{\sqrt{\pi} \alpha M_1},$$

$$c_i = \frac{-\sqrt{\pi} M_1 y_{i+1} + \sqrt{\pi} M_1 y_i + 2\sqrt{h} M_1 y_i^{(\frac{1}{2})} + (\alpha - \beta) D_i + \delta D_{i+1}}{\alpha}$$

$$d_i = \frac{\sqrt{\pi} M_1 M_3 y_{i+1} - \sqrt{\pi} M_1 M_3 y_i - 2\sqrt{h} M_1 M_3 y_i^{(\frac{1}{2})} + (\beta M_3 + \alpha) D_i - (\delta M_3 + \alpha) D_{i+1}}{\alpha M_1}$$

$$e_i = \frac{\sqrt{\pi} M_1 y_{i+1} - \sqrt{\pi} M_1 y_i - 2\sqrt{h} M_1 y_i^{(\frac{1}{2})} + \beta D_i - \delta D_{i+1}}{k\alpha}$$

Such that $\theta = kh$, $M_1 = k \sin \theta$, $M_2 = \cos \theta + \sin \theta - 1$, $M_3 = \cos \theta - 1$,

$$\alpha = -2\sqrt{\pi} M_3 - \sqrt{2\theta} M_2 - \sqrt{\pi h} M_1, \beta = -\sqrt{\pi} M_3 + \sqrt{2\theta} - \sqrt{\pi h} M_1, \delta = -\sqrt{\pi} M_3 + \sqrt{2\theta}, i = 0, 1, 2, \dots, n.$$

We obtain

$$S(x) = \frac{-\sqrt{\pi} M_1 M_3 y_{i+1} + \sqrt{\pi} M_1 M_3 y_i + 2\sqrt{h} M_1 M_3 y_i^{(\frac{1}{2})}}{\alpha M_1} -$$

$$+ \frac{-\sqrt{2k\pi} M_1 M_2 y_{i+1} + \sqrt{2k\pi} M_1 M_2 y_i + 2 M_1 (\alpha + \sqrt{2\theta} l)}{\sqrt{\pi} \alpha} -$$

$$- x_i)^{\frac{1}{2}} + \frac{-\sqrt{\pi} M_1 y_{i+1} + \sqrt{\pi} M_1 y_i + 2\sqrt{h} M_1 y_i^{(\frac{1}{2})} + (\alpha - \beta) D}{\alpha} (6)$$

$$- x_i) + \frac{\sqrt{\pi} M_1 M_3 y_{i+1} - \sqrt{\pi} M_1 M_3 y_i - 2\sqrt{h} M_1 M_3 y_i^{(\frac{1}{2})} + (\beta M_3 + \alpha) D}{\alpha M_1} -$$

$$- x_i) + \frac{\sqrt{\pi} M_1 y_{i+1} - \sqrt{\pi} M_1 y_i - 2\sqrt{h} M_1 y_i^{(\frac{1}{2})} + \beta D_i - \delta D_{i+1}}{k\alpha} \text{sink}(x - x_i).$$

Now apply the fractional continuity condition of the spline function $S_i(x)$ where the splines, $S_{i-1}^{(m)}(x) = S_i^{(m)}(x)$, $m = \frac{1}{2}, 1$, we obtain the following equations

$$D^{(\frac{1}{2})} S_i(x) = D^{(\frac{1}{2})} y_i(x) \quad (7)$$

$$s_{i-1}^{(\frac{1}{2})}(x) = \frac{\sqrt{k}}{2} \left(\frac{-\sqrt{2k\pi} M_1 M_2 y_i + \sqrt{2k\pi} M_1 M_2 y_{i-1}}{\sqrt{\pi} \alpha M_1} + \frac{\sqrt{k}}{2} \left(\frac{-\sqrt{2k}(\beta M_2 + \alpha) D_{i-1} + \sqrt{2k}(\delta M_2 + \alpha) D_i}{\sqrt{\pi} \alpha M_1} \right) \right) + \frac{\sqrt{k}}{2} \left(\frac{\sqrt{\pi} M_1 M_3 y_i - \sqrt{\pi} M_1 M_3 y_{i-1} - 2\sqrt{h} M_1 M_3 y_{i-1}^{(\frac{1}{2})}}{\alpha M_1} \right) (8)$$

$$+ \frac{\sqrt{k}}{2} \left(\frac{\sqrt{\pi} M_1 y_i - \sqrt{\pi} M_1 y_{i-1} - 2\sqrt{h} M_1 y_{i-1}^{(\frac{1}{2})} + \beta D_{i-1} - \delta D_i}{k\alpha} \right)$$

From here by equating equations, (7) and (8) we obtain,

$$A_1 y_i + A_2 y_{i-1} + A_3 y_{i-1}^{(\alpha)} + A_4 D_{i-1} + A_5 D_i = 2k\sqrt{\pi} \alpha M_1 y_i^{(\alpha)} \quad (9)$$

From equation (1),

$$D^y y_i = -p_i(x) y_i' - q_i(x) y_i + r_i(x),$$

$$D^y y_{i-1} = -p_{i-1}(x) y_{i-1}' - q_{i-1}(x) y_{i-1} + r_{i-1}(x), \quad (10)$$

$$y_i' = \frac{y_{i+1} - y_{i-1}}{2h}, \text{ and}$$

$$y_{i-1} = \frac{-y_{i+1} + 4y_i - 3y_{i-1}}{2h}$$

Substitute equation (10) in equation (9) to obtain the iterative formula as finite difference equation

$$a_i y_{i-1} + b_i y_i + c_i y_{i+1} = F_i \quad (11)$$

Then the system of linear equation is formulated from equation (11) as follows

$$Ay = F \quad (12)$$

Where

$$A = \begin{bmatrix} b_1 & c_1 & \dots & \dots & 0 & 0 \\ a_2 & b_2 & c_2 & 0 & 0 & \vdots \\ 0 & a_3 & b_3 & c_3 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & a_{n-1} & b_{n-1} & c_{n-1} \\ 0 & 0 & \dots & 0 & a_n & b_n \end{bmatrix}, y =$$

$$[y_1 \ y_2 \ y_3 \ \dots \ y_{n-1} \ y_n]^T, F = [F - a_1 y_0 \ F_2 \ \dots \ F_{n-1} \ F_n - c_n y_{n+1}]^T.$$

Such that

$$a_i = A_2 + \left(\frac{3p_{i-1}}{2h} - q_{i-1} \right) A_3 - \frac{3A_4 + A_5}{2h} - \frac{\sqrt{\pi k \alpha M_1 p_i}}{h}$$

$$b_i = A_1 - \frac{2A_3 p_{i-1}}{h} + \frac{2A_4}{h} + 2\sqrt{\pi k \alpha M_1} q_i,$$

$$c_i = \frac{A_3 p_{i-1}}{2h} + \frac{A_5 - A_4}{2h} + \frac{\sqrt{\pi k \alpha M_1} p_i}{h},$$

$$F_i = 2\sqrt{\pi k \alpha M_1} r_i(x_i) - A_3 r_{i-1}(x_i), \quad i = 1, 2, \dots, n.$$

$$A_1 = -\sqrt{2} k^{\frac{3}{2}} \pi M_1 M_2 - 4\sqrt{\pi h} k M_1^2 +$$

$$2k^{\frac{3}{2}} \pi M_1 M_3 \cos\left(\theta + \frac{\pi}{4}\right) + 2\pi\sqrt{k} M_1^2 \sin\left(\theta + \frac{\pi}{4}\right),$$

$$A_2 = \sqrt{2} k^{\frac{3}{2}} \pi M_1 M_2 + 4\sqrt{\pi h} k M_1^2 - 2k^{\frac{3}{2}} \pi M_1 M_3 \cos\left(\theta + \frac{\pi}{4}\right) - 2\pi\sqrt{k} M_1^2 \sin\left(\theta + \frac{\pi}{4}\right),$$

$$A_3 = 2k\sqrt{\pi} M_1 (\alpha + \sqrt{2\theta} M_2) + 8\theta M_1^2 - 4\sqrt{\pi \theta} k M_1 M_3 \cos\left(\theta + \frac{\pi}{4}\right) - 2\sqrt{\theta \pi} M_1^2 \sin\left(\theta + \frac{\pi}{4}\right),$$

$$A_4 = -k^{\frac{3}{2}} \sqrt{2\pi} (\alpha + \beta M_2) + 4k\sqrt{h} M_1 (\alpha - \beta) + 2\sqrt{\pi} k^{\frac{3}{2}} (\alpha + \beta M_3) \cos\left(\theta + \frac{\pi}{4}\right) + 2\sqrt{k\pi} M_1 \beta \sin\left(\theta + \frac{\pi}{4}\right),$$

$$A_5 = k^{\frac{3}{2}} \sqrt{2\pi} (\alpha + \delta M_2) + 4k\sqrt{h} M_1 \delta - 2\sqrt{\pi} k^{\frac{3}{2}} (\alpha + \delta M_3) \cos\left(\theta + \frac{\pi}{4}\right) - 2\sqrt{k\pi} M_1 \delta \sin\left(\theta + \frac{\pi}{4}\right).$$

Theorem 3.1: If A is diagonally dominate then for any initial vector $x^{(0)}$ the sequence generates by Gauss-Seidel method $\{x^{(i)}\}_{i=0}^{\infty}$ convergence to unique solution of $Ax = b$.

Proof: see [24].

Theorem 3.2: If A has n independent columns then A is non-singular, A^{-1} exists, and $Au = f$ has a unique solution u .

Proof: see [26] and [21].

Theorem 3.3.[23] Let $\rho \in (0, 1]$, $p \in N$, and $f(x)$ be a continuous function in $[a, b]$ satisfying the following conditions

(i) $D^{j\rho} f(x) \in C([a, b])$ and $D^\rho f(x) \in I([a, b])$, $\forall j = 1, 2, \dots, p$.

(ii) $D^{(p+1)\rho} f(x)$ is continuous on $[a, b]$.

Then for each $x \in [a, b]$,

$$f(x) = \sum_{j=0}^p \frac{D^{j\rho} f(a)(x-a)^{j\rho}}{\Gamma(j\rho+1)} + R_p(x, a), \text{ where}$$

$$R_p(x, a) = D^{(p+1)\rho} f(\eta) \frac{(x-a)^{(p+1)\rho}}{\Gamma((p+1)\rho+1)}, \quad a \leq \eta \leq x.$$

4. Error Estimations and Convergence Analysis

Theorem 4.1: Let $S(t)$ be the unique non-polynomial fractional spline satisfying in (4), for a given function $y(x) \in C^5[a, b]$. Then the following error estimates hold:

$$|s^{(m)}(x) - f^{(m)}(x)| \leq \frac{h^{m+\alpha}}{(2m-2)!} \|f^{(6)}\|, \quad \text{where } \alpha \in R$$

and

$$|e^{(m)}(x)| = |f^{(m)}(x) - S^{(m)}(x)|, \quad \text{and, } w = \max_{0 \leq x \leq h} |f^{(2n)}(x)|.$$

Proof: By subtracting the analytic function $y(x)$ of sufficiently high order with the spline model in equation (6) and using theorem 3.3, we obtain.

Since $D^{\frac{1}{2}}s(x)$ is Hermite interpolation polynomial of degree 3, and matching $D^{\frac{1}{2}}f(x), D^{\frac{3}{2}}f(x)$, at $x = x_j, x_{j+1}$ so for any $x \in [x_j, x_{j+1}]$, using ([27],[28]) and let

$m = 3, g = f^{(\frac{1}{2})}$ and $p_3 = s^{(\frac{3}{2})}(x)$, form the non-polynomial spline function in equation (4), with known constraint conditions; we get:

$$\left| D^{\frac{1}{2}}s(x) - D^{\frac{1}{2}}f(x) \right| \leq \frac{h^5}{4!} |D^{(6)}f(x)|, \text{ also if we put}$$

$g = D^{\frac{3}{2}}f(x)$ and $P_3 = D^{\frac{3}{2}}s(x)$, we get

$$\left| s^{(\frac{3}{2})}(x) - f^{(\frac{3}{2})}(x) \right| \leq \frac{h^3}{3!} |D^{(6)}f(x)|, \text{ then}$$

$$|s(x) - s(0) + f(0) - f(x)| \leq \frac{h^6}{6!} \|f^{(6)}(x)\|$$

Since $s(0) = f(0)$ and $x \in [0, 1]$ then the last equation becomes

$$\left| s^{(\frac{5}{2})}(x) - f^{(\frac{5}{2})}(x) \right| \leq \frac{h^{\frac{7}{2}}}{2!} \|f^{(6)}(x)\|,$$

and since $f^{(p)}(0) = 0, p = 1, 2$, also using [23], clearly find the error estimation as follows following:

- i. let $\zeta = 0, m = 4$, then $|e(x)| \leq \frac{h^8}{8!} \|f^{(8)}(x)\|_{\infty}$.
- ii. let $\zeta = 1, m = 4$, then $|e^{(\frac{1}{2})}(x)| \leq \frac{h^7}{6!} \|f^{(8)}(x)\|_{\infty}$.
- iii. let $\zeta = 1, m = 4$, then $|e'(x)| \leq \frac{h^6}{24!} \|f^{(8)}(x)\|_{\infty}$.
- iv. let $\zeta = \frac{3}{2}, m = 4$, then $|e^{(\frac{3}{2})}(x)| \leq \frac{h^5}{32!} \|f^{(8)}(x)\|_{\infty}$.
- v. let $\zeta = 2, m = 4$, then $|e^{(2)}(x)| \leq \frac{h^4}{4!} \|f^{(8)}(x)\|_{\infty}$.

5. Numerical experiments

In this section the method applied to solve two numerical examples of boundary fractional differential equations with constant coefficients, the result compared with the exact analytical solution to show the efficiency of the method. The computational programs were written in MatLab. Here the algorithms of the Gauss-Seidel and the conjugate gradient methods are presented.

Algorithm 5.1

Suppose that we have the linear system (12) where A is symmetric positive definite matrix for Gauss-Seidel method first decompose matrix A as $A = D + L + U$ such that D is diagonal matrix, L is lower matrix and U is upper matrix. Then the Gauss-Seidel algorithm can be written as:

Start with initial vector $y^{(0)}$.

$$y^{(i+1)} = -(D + L)^{-1}Uy^{(i)} + (D + L)^{-1}F, i = 0, 1, 2, \dots$$

Algorithm 5.2 (conjugate gradient method)

Chose $y_0 \in R^n$ and put, $d_0 = r_0 = F - Ay_0$.

For $k = 0, 1, 2, \dots$

If $d_k = 0$, stop y_k is solution of $Ay = F$.

Otherwise Compute

$$\alpha_k = \frac{r_k^t r_k}{d_k^t A d_k}, y_{k+1} = y_k + \alpha_k d_k,$$

$$r_{k+1} = r_k - \alpha_k A d_k, \beta_k = \frac{r_{k+1}^t r_{k+1}}{r_k^t r_k},$$

$$d_{k+1} = r_{k+1} + \beta_k d_k$$

Example 5.1. Consider the fractional differential equation

$$y' + y^\alpha + y = 2\sin 2x + 3\cos 2x, \quad x \in [0, \pi], y(0) = y(\pi) = 0 \tag{13}$$

The exact solution of (13) when $\alpha = \frac{1}{2}$ is, $y = \sin 2x$.

Table 1 shows the number of iterations with different value of j using Gauss-Seidel and Conjugate gradient methods.

Table 1. Iteration numbers of example 5.1.

Number of iterations			
j	64	128	256
GS	161	687	6962
CG	63	127	255

Table 2 Exact, approximate, and absolute error of example 5.1.

x	Exact	Approximate	Absolute error
---	-------	-------------	----------------

	solution	solution	
$\frac{\pi}{64}$	0.098	0.0973670	6.5×10^{-4}
$\frac{\pi}{8}$	0.7071	0.7031279	3.9×10^{-3}
$\frac{\pi}{4}$	1	0.9957816	4.2×10^{-3}
$\frac{3\pi}{8}$	0.7071	0.7062436	8.6×10^{-4}
$\frac{\pi}{2}$	0	0.0038903	3.8×10^{-3}




Figure 1 Exact and approximate solution of example 5.1 with $h = \frac{\pi}{64}$




Figure 2 Exact and approximate solution of example 5.1 with $h = \frac{\pi}{128}$

Example 5.2. A boundary value problem of FDE

$$y' + y = x^2 + x + \frac{8}{3}\sqrt{\frac{x^3}{\pi}} - 2\sqrt{\frac{x}{\pi}} - 1 + y^\alpha, x \in [0, 1], y(0) = 0, \text{ and } y(1) = 0.$$

The exact solution with $\alpha = \frac{1}{2}$ is given as, $y = x^2 - x$.

The iteration numbers is present in Table 3

Table 3 iteration numbers of example 5.2.

j	Number of iterations		
	64	128	256
GS	220	1607	7838
CG	63	127	255

Table 4 Exact, approximate, and absolute error of example 5.2.

x	Exact solution	Approximate solution	Absolute error
$\frac{1}{64}$	-0.01538	-0.0134754	1.9×10^{-3}
$\frac{1}{8}$	-0.1093	-0.0982452	1.1×10^{-2}
$\frac{1}{4}$	-0.1875	-0.1772538	1.7×10^{-2}

$\frac{3}{8}$	-0.2343	-0.23071	3.6×10^{-3}
$\frac{1}{2}$	-0.25	-0.254126	4.1×10^{-3}




Figure 3 exact and approximate solution of example 5.2 $h = \frac{1}{128}$




Figure 4 exact and approximate solution of example 5.2 $h = \frac{1}{256}$

Example 5.3.[29] Consider the fractional differential equation

$$D^\alpha y(x) = -y(x) + x^4 - \frac{1}{2}x^3 - \frac{3}{\Gamma(4-\alpha)}x^{3-\alpha} + \frac{4}{\Gamma(5-\alpha)}x^{4-\alpha}, 0 < \alpha < 1.$$

With boundary conditions, $y(0) = 0, y(1) = \frac{1}{2}$, and $x \in [0,1]$.

The exact solution with $\alpha = \frac{1}{2}$ is, $y = x^4 - \frac{1}{2}x^3$.

Table 5. Iteration numbers of example 5.3

Number of iterations	
J	1000
CG	999
GS	19710

Table 6. Exact, approximate, and absolute error of example 5.3.

x	Exact solution	Approximate solution	Absolute error
0.1	-0.0004	-0.00011694	2.8×10^{-4}
0.2	-0.0024	-0.000194	2.2×10^{-3}
0.3	-0.0054	-0.00003	5.4×10^{-3}

0.4	-0.0064	0.00140836	7.8×10^{-3}
0.5	0	0.00647904	6.4×10^{-3}
0.6	0.0216	0.02230459	7.0×10^{-4}
0.7	0.0686	0.06383659	4.7×10^{-3}
0.8	0.1536	0.15236168	1.2×10^{-3}
0.9	0.2916	0.30210056	1.05×10^{-2}

6. Conclusion

This paper developed the trigonometric spline method for solving FDE and conformable with conjugate gradient methods. The findings with the non-polynomial quartic spline functions are really quite interesting. In approximating functions, the non-polynomial spline and conjugate gradient approaches are more adaptive, as seen in the numerical examples. The graphs comparing exact and approximate solutions for numerical examples demonstrate our method's superiority.

7. References

- [1]. Torvik, P.J. and R.L. Bagley, *On the appearance of the fractional derivative in the behavior of real materials*. 1984.
- [2]. Engheta, N., *On fractional calculus and fractional multipoles in electromagnetism*. IEEE Transactions on Antennas and Propagation, 1996. **44**(4): p. 554-566.
- [3]. Assaleh, K. and W.M. Ahmad. *Modeling of speech signals using fractional calculus*. in *2007 9th International Symposium on Signal Processing and Its Applications*. 2007. IEEE.
- [4]. Gafiychuk, V., B. Datsko, and V. Meleshko, *Mathematical modeling of time fractional reaction-diffusion systems*. Journal of Computational and Applied Mathematics, 2008. **220**(1-2): p. 215-225.
- [5]. Carpinteri, A. and F. Mainardi, *Fractals and fractional calculus in continuum mechanics*. Vol. 378. 2014: Springer.
- [6]. Ali, M.F., M. Sharma, and R. Jain, *An application of fractional calculus in electrical engineering*. Adv. Eng. Tec. Appl, 2016. **5**(4): p. 41-45.
- [7]. Tarasov, V.E., *On history of mathematical economics: Application of fractional calculus*. Mathematics, 2019. **7**(6): p. 509.

- [8]. Oldham, K. and J. Spanier, *The fractional calculus theory and applications of differentiation and integration to arbitrary order*. 1974: Elsevier.
- [9]. Jafari, H., M. Dehghan, and K. Sayevand, *Solving a fourth-order fractional diffusion-wave equation in a bounded domain by decomposition method*. Numerical Methods for Partial Differential Equations: An International Journal, 2008. **24**(4): p. 1115-1126.
- [10]. Li, C. and C. Tao, *On the fractional Adams method*. Computers & Mathematics with Applications, 2009. **58**(8): p. 1573-1588.
- [11]. Hashim, I., O. Abdulaziz, and S. Momani, *Homotopy analysis method for fractional IVPs*. Communications in Nonlinear Science and Numerical Simulation, 2009. **14**(3): p. 674-684.
- [12]. Daraghme, A., N. Qatanani, and A. Saadeh, *Numerical solution of fractional differential equations*. Applied Mathematics, 2020. **11**(11): p. 1100-1115.
- [13]. Zabidi, N.A., et al., *Numerical Solutions of Fractional Differential Equations by Using Fractional Explicit Adams Method*. Mathematics, 2020. **8**(10): p. 1675.
- [14]. Bhatti, M.I. and M. Rahman, *Technique to Solve Linear Fractional Differential Equations Using B-Polynomials Bases*. Fractal and Fractional, 2021. **5**(4): p. 208.
- [15]. Mohammed, P.O., T. Abdeljawad, and F.K. Hamasalh, *Discrete Prabhakar fractional difference and sum operators*. Chaos, Solitons & Fractals, 2021. **150**: p. 111182.
- [16]. Srivastava, H.M., et al., *Solutions of General Fractional-Order Differential Equations by Using the Spectral Tau Method*. Fractal and Fractional, 2022. **6**(1): p. 7.
- [17]. Wright, S. and J. Nocedal, *Numerical optimization*. Springer Science, 1999. **35**(67-68): p. 7.
- [18]. Fletcher, R. and C.M. Reeves, *Function minimization by conjugate gradients*. The computer journal, 1964. **7**(2): p. 149-154.
- [19]. Ivanov, B., et al., *A Novel Value for the Parameter in the Dai-Liao-Type Conjugate Gradient Method*. Journal of Function Spaces, 2021. **2021**.
- [20]. Justine, H. and J. Sulaiman, *Cubic non-polynomial solution for solving two-point boundary value problems using SOR iterative method*. Transactions on Science and Technology, 2016. **3**(3): p. 469-475.
- [21]. Hamasalh, F.K. and M.A. Headayat. *The applications of non-polynomial spline to the numerical solution for fractional differential equations*. in *AIP Conference Proceedings*. 2021. AIP Publishing LLC.
- [22]. Al Bayati, A.Y., R.K. Saeed, and F.K. Hama-Salh, *The Existence, Uniqueness and Error Bounds of Approximation Splines Interpolation for Solving Second-Order Initial Value Problems I*. 2009.
- [23]. Hamasalh, F.K. and P.O. Muhammad, *Numerical Solution of Fractional Differential Equations by using Fractional Spline Functions*. Journal of Zankoy Sulaimani - Part A, 2015. **17**: p. 97-110.
- [24]. Burden, R.L., J.D. Faires, and A.M. Burden, *Numerical analysis*. 2015: Cengage learning.
- [25]. Bulirsch, R., J. Stoer, and J. Stoer, *Introduction to numerical analysis*. Vol. 3. 2002: Springer.
- [26]. Yano, M., et al., *Math, Numerics, & Programming (for Mechanical Engineers)*. 2012.
- [27]. Birkhoff, G. and A. Priver, *Hermite interpolation errors for derivatives*. Journal of Mathematics and Physics, 1967. **46**(1-4): p. 440-447.
- [28]. Hamasalh, F.K. and A.H. Ali. *On the generalized fractional cubic spline with application*. in *AIP Conference Proceedings*. 2019. AIP Publishing LLC.
- [29]. Al-Rabtah, A., S. Momani, and M.A. Ramadan. *Solving linear and nonlinear fractional differential equations using spline functions*. in *Abstract and Applied Analysis*. 2012. Hindawi.

الدالة سبلاين المثلثية المحسنة مع طريقة التدرج المقترن لحل FDEs

فريدون قادر حمه صالح ، غولنار وسيم صادق ، عماد صنعان سلام^٢

^١قسم الرياضيات، كلية التربية، جامعة السليمانية، السليمانية، اقليم كردستان-العراق

^٢قسم الرياضيات، كلية التربية الأساس، جامعة السليمانية، السليمانية، اقليم كردستان-العراق

^٣قسم الرياضيات، كلية التربية، جامعة السليمانية، السليمانية، اقليم كردستان-العراق

الخلاصة:

نحن نبحت في دالة سبلاين غير متعددة الحدود لحل المعادلات التفاضلية الكسرية باستخدام طريقة التدرج المترافق المطابق. تم وصف المشتق الكسري واستخدام التكامل الجزئي والمشتق (Caputo) يجب أن يبني الاستيفاء الخطي باستخدام معاملات متعددة الحدود الكسري. هذا ، بالتالي ، يحول المشكلة إلى نظام خطي تكراري مكافئ يمكن حله بواسطة (Gauss-Seidel) وطرق التدرج المترافق. بالنسبة لوظيفة الشريحة المحددة ، تمت دراسة حدود الخطأ واكتمل تحليل الثبات ، ويتم حساب تقدير الخطأ أيضًا نظرًا لأن قيم مختلفة لـ (n) تعتمد على حجم الخطوة (h) . تم عرض أمثلة عددية مع حلول تحليلية معروفة للتحقق من دقة الطريقة. كانت النتائج في علاقة مرضية مع الإجابات الدقيقة وفقًا للتجارب العددية. علاوة على ذلك ، تم التحقيق في تحليل التقارب مع اشتقاق بعض النظريات . أيضًا ، يتم شرح الإجراء بعمق ودعمه بأمثلة حسابية وتظهر النتائج أن وظيفة الشريحة الجزئية التي تقم البيانات منتجة ومربحة في حل المشكلات الفريدة ومقارنتها بالحلول الدقيقة.