

Supra semi preopen sets in supra topological spaces

N. A. Nadhim¹, Ahmed Hanoon Abud²

¹Department of Mathematics, Faculty of Education for pure science, University of AL-Anbar, AL-Anbar, Iraq;

²Department of Mathematics, College of Science, University of AL-Mustnsiriyah, Baghdad, Iraq;



ARTICLE INFO

Received: 30 / 10 / 2023

Accepted: 26 / 11 / 2023


Available online: 19 / 12 / 2023

DOI: 10.37652/juaps.2023.181587

Keywords:

supra topological space, preopen set, supra preopen set, supra semi preopen set, supra pre compact space.

Copyright©Authors, 2022, College of Sciences, University of Anbar. This is an open-access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).



ABSTRACT

The purpose of this article is to provide a new set which is supra semi preopen set via supra topological space (X, \mathcal{M}_X) . with the help of some examples and properties. Also, some of their properties have been investigate. With we illustrate the relationships between this concept and supra preopen set (respectively, supra open set). Also, we define two spaces which are supra semi pre compact and supra semi pre Lindelöf spaces with the relationships between them. At the end of this work, we provide some examples, properties to support this work.

1-Introduction

In 1983, Mashhour [1] introduced the concept of supra topological space, which is, for any set $X \neq \emptyset$, and the collection M_X of subset of X that \emptyset and X belong to M_X , also arbitrary union of elements of M_X is an element in M_X . The pair (X, M_X) is called supra topological space (briefly, su.top.sp), and the elements of M_X are said to be supra open (briefly, su.o) sets and its complements are supra closed (briefly, su.c) sets. Also, he presented the relation between this concept and topological space (for short top. sp.) which is (Every top.sp is su.top.sp). The auther in [2], the concept of supra interior was defined, the supra interior of a subset \mathcal{A} of supra space (X, M_X) , which is $su.int(\mathcal{A}) = \bigcup \{U : U \subseteq \mathcal{A}, \text{ where } U \in M_X\}$, and supra closure of \mathcal{A} , that is $su.cl(\mathcal{A}) = \bigcap \{F : \mathcal{A} \subseteq F, \text{ where } F^c \in M_X\}$,

After then in 2010, Sayed [3] provided a new concept which is supra preopen (respectively, supra pre closed) set, briefly, su. pr.o (respectively, su. pr. c) set, such that \mathcal{A} is su.pr.o if $\mathcal{A} \subseteq su.int(su.cl(\mathcal{A}))$.

*Corresponding author at: Department of Mathematics, Faculty of Education for pure science, University of AL-Anbar, AL-Anbar, Iraq; Iraq; ORCID: <https://orcid.org/0000-0000-0000-0000>; Tel:+9647805269952 E-mail address: mad772918@uoanbar.edu.iq

The complement of a su. pr. o set is said to be su.pr.c, The collection of all su. pr. o (respectively, su.pr.c) sets of X is denoted by $su.pr.o(X)$ (respectively, $su. pr.c(X)$). And this author defined, The union of all $su.pr.o(X)$ set contained in a subset \mathcal{A} of su.sp. (X, M_X) is su.pr.interior of \mathcal{A} and we denoted by $su.int_p(\mathcal{A})$. Also, The intersection of all $su.pr.c(X)$ sets containing \mathcal{A} is said to be su.pr closure of \mathcal{A} and we denoted by $su.cl_p(\mathcal{A})$. It is well known that top. sp and su. top. sp have been generalized and studied in many ways (see for example, [4], [5], [6]).

In 2020, Al-Shami and others [7], presented the concept of supra pre compact and supra pre Lindelöf spaces with many properties, examples and theorem with the relationships between these two spaces.

In this work, we provide a new concept which is supra semi* preopen (briefly, su. s* pr.o) set in su.top.sp and the fact that is "every su.o set is su. s* pr.o set", but the convers is not true, also the relation between su. pr.o set and su. s* pr.o set, see (Remark 2.2, part(2)) in this research, Finally, we provide anew spaces namely su. s* pr. compact and su. s* pr. Lindelöf spaces (briefly, su. s* pr.com and su. s* pr. Lin) with some properties, examples for these spaces.

2- supra. s*pre. open set

At 2002, Veera Kumar [8], introduce the concept semi- pre-open set. By the same context via supra topology, we define the following: -

Definition 2. 1: - A subset \mathcal{A} of su. sp $(\mathcal{X}, M_{\mathcal{X}})$ is said to be su. s*pr. o, if there exists su. pr.o set. \mathcal{G} , such that $\mathcal{G} \subseteq \mathcal{A} \subseteq \text{su.cl}_p(\mathcal{A})$

Remarks 2. 2: -

1-Every su. o set. is su. s*pr. o set. but the converse may be not true.

2- Every su. pr.o set is su.s*pr.o set , but the convers is not true.

Example 2. 3: - In su. indiscrete sp. (\mathcal{R}, M_{ind}) , the set of all rational numbers is su. s*pr. o set. but not su.o set.

Example 2. 4: - Let $\mathcal{X} = \{1,2,3\}, M_{\mathcal{X}} = \{\emptyset, \mathcal{X}, \{1\}, \{3\}, \{1,3\}, \{2,3\}\}$.

Then, $M_{\mathcal{X}}^c = \{\emptyset, \mathcal{X}, \{1\}, \{2\}, \{1,2\}, \{2,3\}\}$

the su. pr. o $(\mathcal{X}) = \{\emptyset, \mathcal{X}, \{1\}, \{3\}, \{1,3\}, \{2,3\}\}$

and su.s*pr. o $(\mathcal{X}) = \{\emptyset, \mathcal{X}, \{1\}, \{3\}, \{1,3\}, \{2,3\}, \{1,2\}\}$,

we note that $\{1,2\}$ is su.s*pr. o set but not su. pr. o set.

Proposition 2. 5: - The union of any family of su. s*pr.o (\mathcal{X}) is su. s*pr. o (\mathcal{X})

Proof: - Let $\{A_{\alpha}\}_{\alpha \in \Omega}$ be a family of su. s*pr. o set of \mathcal{X} , so there exists pr.o subset $\{U_{\alpha}\}_{\alpha \in \Omega}$ such that $U_{\alpha} \subseteq A_{\alpha} \subseteq \text{cl}_p(U_{\alpha})$, for each $\alpha \in \Omega$.

Also, from (theorem2 [3]) $\bigcup_{\alpha \in \Omega} U_{\alpha} \subseteq \bigcup_{\alpha \in \Omega} A_{\alpha} \subseteq \bigcup_{\alpha \in \Omega} \text{su.cl}(U_{\alpha}) \subseteq \text{su.cl}(\bigcup_{\alpha \in \Omega} U_{\alpha})$

Then, $\bigcup_{\alpha \in \Omega} A_{\alpha}$ is su. s*pr. o set.

Corollary 2. 6: - The intersection of any family of su. s*pr.c (\mathcal{X}) is su. s*pr.c (\mathcal{X}) .

Proof:- Let $\{F_{\alpha}\}_{\alpha \in \Omega}$ be su.s*pr.o sets, by (theorem2 [3]), we get $\bigcup_{\alpha \in \Omega} \{\mathcal{X} - F_{\alpha}\}$ be su.s*pr.o set, but $\bigcup_{\alpha \in \Omega} \{\mathcal{X} - F_{\alpha}\} = \mathcal{X} - \bigcap_{\alpha \in \Omega} \{F_{\alpha}\}$,

so $\mathcal{X} - (\mathcal{X} - \bigcap_{\alpha \in \Omega} \{F_{\alpha}\}) = \bigcap_{\alpha \in \Omega} \{F_{\alpha}\}$ is su. s*pr.cl (\mathcal{X}) .

Remark 2. 7: - The intersection of two su. s*pr.o sets is may be not su. s*pr.o set.

Example 2. 8: In (Example2.4) the intersection of two su.s*pr.o sets $\{1,2\} \cap \{2,3\} = \{2\}$ is not su.s*pr.o set.

Proof: - Suppose \mathcal{A} is su. pr.o set and $\mathcal{A} \subseteq \mathcal{A} \subseteq \text{su.cl}_p(\mathcal{A})$, then it is su.s*pr.o set .

Proposition 2. 9: - A subset F of su.sp $(\mathcal{X}, M_{\mathcal{X}})$ is su.s*pr.c set if and only if $F = \text{su.s*pr.cl}(F)$.

Proof: - Let F be su.s*pr.c set, the smallest intersection of all su.s*pr.c sets which contain F is equal to F. Therefore, $F = \text{su. s*pr.cl}(F)$.

Conversely, since $\text{su.s*pr.cl}(F)$ always su.s*pr.c set, by corollary (2.6), Then, F is also su.s*pr.c set.

Corollary 2. 10: - For a subset F of su.sp $(\mathcal{X}, M_{\mathcal{X}})$, we have $\text{su.s*pr.cl}(\text{su.s*pr.cl}(F)) = \text{su.s*pr.cl}(F)$

Proof: - Since by corollary (2.6), $\text{su.s*pr.cl}(F)$ is su.s*pr.c set, then by (proposition 2.9), we get the result.

Definition 2. 11: - A point $x \in \mathcal{X}$, is said to be su.s*pr. interior point to a subset \mathcal{A} of su.sp $(\mathcal{X}, M_{\mathcal{X}})$, if there is su.s*pr.o set U with $x \in U \subseteq \mathcal{A}$.

Definition 2. 12: - A point $x \in \mathcal{X}$, is said to be su.s*pr-adherent point to a subset F of su.sp $(\mathcal{X}, M_{\mathcal{X}})$, if $U \cap F \neq \emptyset$, where U is su.s*pr.o set containing x .

Proposition 2. 13: -For a subset G of su.sp $(\mathcal{X}, M_{\mathcal{X}})$ we have $\text{su.s*pr.int}(\mathcal{X} - G) = \mathcal{X} - (\text{su.s*pr.cl}(G))$.

Proof:- Suppose $x \notin \mathcal{X} - (\text{su.s*pr.cl}(G))$, so $x \in \text{su.s*pr.cl}(G)$, then for each su.s*pr.o set U containing x , we get $U \cap G \neq \emptyset$, that is x is not su.s*pr. interior point to $\mathcal{X} - G$, that is $x \notin \text{su.s*pr.int}(\mathcal{X} - G)$, then $\text{su.s*pr.int}(\mathcal{X} - G) \subseteq \mathcal{X} - (\text{su.s*pr.cl}(G))$

Conversely, Let $x \notin \text{su.s*pr.int}(\mathcal{X} - G)$, then there is su.s*pr.o set U containing x with $U \not\subseteq \mathcal{X} - G$, then $U \cap G \neq \emptyset$, so x is su.s*pr-adherent point to G , that is $x \in \text{su.s*pr.cl}(G)$. Hence, $x \notin \mathcal{X} - (\text{su.s*pr.cl}(G))$, then $\mathcal{X} - (\text{su.s*pr.cl}(G)) \subseteq \text{su.s*pr.int}(G)$. Hence, $\text{su.s*pr.int}(\mathcal{X} - G) = \mathcal{X} - (\text{su.s*pr.cl}(G))$.

Proposition 2. 14: - A subset G of su.sp $(\mathcal{X}, M_{\mathcal{X}})$ is su.s*pr.o set if and only if $\text{su.s*pr.int}(G) = G$.

Proof: - Let $\text{su.s*pr.int}(G) = G$, but the union of su.s*pr.o set is also su.s*pr.o set, then $\text{su.s*pr.int}(G)$ is su.s*pr.o set, which equal to G , then G is su.s*pr.o set.

Conversely, suppose G is su.s*pr.o set, so it is su.s*pr-neighborhood of each its point, so $\text{su.s*pr.int}(G) = G$.

3- su.s*pr - Lindelöf space

In this section, we introduce new concepts namely su.s*pr-compact space (briefly, su.s*pr-com. sp) and su.s*pr-Lindelöf space (briefly, su.s*pr-Lind. sp).

Definition 3. 1: - A su.sp $(\mathcal{X}, M_{\mathcal{X}})$ is said to be su.s*pr-com.sp , if for each su.s*pr.o cover to \mathcal{X} has a finite subcover.

Definition 3. 2: - A $su.sp(\mathcal{X}, M_{\mathcal{X}})$ is said to be $su.s^*pr$ -Lind. sp, if for each $su.s^*po$ cover to \mathcal{X} has a countable subcover.

Remark 3. 3: - Every $su.s^*pr$ -com sp is $su.s^*pr$ -Lind. sp.

Proposition 3. 4: - Every $su.s^*pr.c$ subset of $su.s^*pr$ -Lind. sp is $su.s^*pr$ -Lind.

Proof: - Let K be $su.s^*pr.c$ subset of $su.s^*pr$ -Lind. sp $(\mathcal{X}, M_{\mathcal{X}})$ and $\mathcal{C} = \{V_{\alpha} : \alpha \in \Omega\}$ be $su.s^*pr.o$ cover to K , that is $K \subseteq \bigcup_{\alpha \in \Omega} V_{\alpha}$, so $\mathcal{X} \subseteq (\bigcup_{\alpha \in \Omega} V_{\alpha}) \cup K^c$, where K^c is $su.s^*pr.o$ set, but \mathcal{X} is $su.s^*pr$ -Lind., then $\mathcal{X} \subseteq (\bigcup_{i=1}^{\infty} V_{\alpha_i}) \cup K^c$. Hence, $\mathcal{X} \subseteq \bigcup_{i=1}^{\infty} V_{\alpha_i}$, then K is $su.s^*pr$ -Lind.

Proposition 3.5: -The union of two $su.s^*pr$ -Lind. subsets of $su.sp(\mathcal{X}, M_{\mathcal{X}})$ is $su.s^*pr$ -Lind

Proof: - Let H and K be two $su.s^*pr$ -Lind. subsets and $\mathcal{C} = \{V_{\alpha} : \alpha \in \Omega\}$ be $su.s^*pr.o$ cover to $H \cup K$, that is, $H \cup K \subseteq \bigcup_{\alpha \in \Omega} V_{\alpha}$, then \mathcal{C} is $su.s^*po$ cover to H and K , then $H \subseteq \bigcup_{i=1}^{\infty} V_{\alpha_i}$ and $K \subseteq \bigcup_{j=1}^{\infty} V_{\alpha_j}$, since H and K are $su.s^*p$ -Lind. set. So, $H \cup K \subseteq \bigcup_{i,j=1}^{\infty} V_{\alpha_{ij}}$.

Hence, $H \cup K$ is $su.s^*p$ -Lind. set.

Proposition 3. 6: -A $su.sp(\mathcal{X}, M_{\mathcal{X}})$ is $su.s^*p$ -comp. if and only if every collection of $su.s^*pc$ set of \mathcal{X} with FIP has non empty intersection.

Proof:- Let \mathcal{X} be $su.s^*p$ -comp. and $\{G_{\alpha} : \alpha \in \Omega\}$ be $su.s^*pc$ subsets of \mathcal{X} with assume that $\bigcap_{\alpha \in \Omega} G_{\alpha} = \emptyset$, then $(\bigcap_{\alpha \in \Omega} G_{\alpha})^c = \emptyset^c$, then $\bigcup_{\alpha \in \Omega} G_{\alpha}^c = \mathcal{X}$, but for each $\alpha \in \Omega$, we have G_{α} is $su.s^*po$ set, so $\{G_{\alpha}^c : \alpha \in \Omega\}$ be a cover of $su.s^*po$ sets of \mathcal{X} , which is $su.s^*p$ -comp., then $\mathcal{X} \subseteq \bigcup_{i=1}^n G_{\alpha_i}^c$, so $\mathcal{X}^c = (\bigcup_{i=1}^n G_{\alpha_i}^c)^c$. Hence, $\emptyset = \bigcap_{i=1}^n G_{\alpha_i}$, which is a contradiction with FIP, therefore $\bigcap_{\alpha \in \Omega} G_{\alpha} \neq \emptyset$.

Proposition 3. 7: - Let $(\mathcal{X}, M_{\mathcal{X}})$ be $su.sp$, then every subspace of \mathcal{X} is $su.s^*p$ -Lind., if every $su.s^*po$ of \mathcal{X} is $su.s^*pr$ -Lind.

Proof: - Let H be any set in \mathcal{X} and $\{S_{\alpha} : \alpha \in \Omega\}$ be $su.s^*po$ cover to H . So, $H \subseteq \bigcup_{\alpha \in \Omega} S_{\alpha}$, by proposition 2.5, $\bigcup_{\alpha \in \Omega} S_{\alpha}$ is $su.s^*po$, hence it is $su.s^*p$ -Lind., then $H \subseteq \bigcup_{\alpha \in \Omega} S_{\alpha} \subseteq \bigcup_{i=1}^{\infty} S_{\alpha_i}$. Therefore, H is $su.s^*p$ -Lind..

Conclusion

In this work, we presented a type of a set, which is Supra semi preopen sets in supra topological spaces, with some characteristics, examples, and theories associated with that sets. Whoever reads this work should consider other groups in the topological space such as Supra ω -preopen set in supra topological spaces or Supra θ -preopen set in supra topological spaces with proceed with the same research method to reach what is desired.

Acknowledgment

In this humble work, I would like to extend my thanks and appreciation to the distinguished professor, Professor Dr. Haider Jebur Ali, for the assistance he provided me in completing this work.

References

- [1] A. S. Mashhour, A. A. Allam, F. S. Mahmoud, F. H. Khedr. (1983). On supra topological space. *Indian Journal of Pure and Applied Mathematics*, 14, 502-510.
- [2] O. R. Sayed. (2012). Supra β -connectedness on topological spaces. *Pakistan Academy of Sciences*, 49, 19-23.
- [3] O. R. Sayed. (2010). Supra pre-open sets and supra pre-continuity on topological spaces, *Scientific Studies and Research*, 20, 79-88.
- [4] Abo-elhamaye and T. M. Al-shami. (2016). Supra homeomorphism in supra topological ordered spaces, *Facts Universitatis, Series: Mathematics and Informatics*, 31 (5) , 1091-1106.
- [5] N. K. Humadi, H. J. Ali. (2019). Supra ωb -separation axioms. *Al-Mustnsiriyah Journal of Science*, 30, 88-95.
- [6] T. M. Al-Shami. (2016). Some results related to supra topological spaces. *Journal of Advanced Studies in Topology*, 7, 283-294.
- [7] T. M. Al-Shami. B. A. Asaad. M. A. El-Gayar. (2020). Various Types of Supra Pre-compact and Supra Pre-Lindelöf Spaces. *Missouri J. Math. Sci.* 32 (1) 1-20, May2020. <https://doi.org/10.35834/2020/3201001>.
- [8] M. K. R. S. Veera Kumar. (2002). Pre semi – closed sets. *Indian Journal of Mathematics*, Vol. 44, No.2,165-181.

المجاميع المفتوحة من النمط شبه الفوقية في الفضاءات الطوبولوجية الفوقية

ناديه علي ناظم^{1*}، احمد حنون عبود²

¹ قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعه الانبار، الانبار، العراق.

² قسم الرياضيات، كلية العلوم، الجامعة المستنصرية، بغداد، العراق

mad772918@uoanbar.edu.iq

ahmedhanoon@uomustansiriyah.edu.iq

الخلاصة:

الغرض من هذا البحث هو تقديم مجموعة جديدة هي المجاميع المفتوحة من النمط شبه الفوقية في الفضاءات الطوبولوجية الفوقية بمساعدة بعض الأمثلة والخصائص. كما تم التحقيق في بعض تلك الخصائص. مع توضيح العلاقات بين هذا المفهوم والمجموعة المفتوحة أعلاه (على التوالي، المجموعة المفتوحة من النمط شبه الفوقية). أيضاً، قمنا بتعريف فضاءين هما الفضاء شبه المرصوص الاعلى وفضاء شبه اللايندولوف الاعلى مع توضيح العلاقات بينهما. وفي نهاية هذا العمل، قدمنا بعض الأمثلة والخصائص لدعم هذا العمل.

الكلمات المفتاحية: - الفضاء الطوبولوجي الفوقي. المجموعة قبل المفتوحة. المجموعة قبل المفتوحة الفوقية. المجموعة قبل المفتوحة شبه الفوقية. الفضاء المرصوص قبل الفوقية.